



GREATER BOSTON INFUSION CENTER

17/19 Kendrick St, Wrentham, MA 02093

P 774-581-0032 | F 774-581-0033

PAGE 1

Benlysta - Referral Checklist

Referral packet checklist

Complete this page and attach the supporting items below.

PATIENT & REFERRING PROVIDER

PATIENT NAME

DOB

DATE

REFERRING PROVIDER

NPI

DIRECT CONTACT / EXT.

OFFICE / PRACTICE

PHONE

FAX

EMAIL

URGENT CLINICAL CONTACT

REQUIRED REFERRAL PACKET

- ☐ Patient demographics and contact information
- ☐ Primary and secondary insurance information; front/back card copies
- ☐ Supporting clinical notes / office visits documenting Benlysta treatment plan
- ☐ Current medication list and allergies
- ☐ Relevant laboratory results / imaging and baseline safety information
- ☐ Completed Benlysta prescribing order (following page)

MEDICATION-SPECIFIC CLINICAL DOCUMENTATION

- Note should include any therapies tried/failed, and must include discussion about Benlysta
- Medication list and allergies
- Recommended: CBC w diff, JC Virus, negative TB screening

GBIC INTERNAL USE

REFERRAL RECEIVED

BENEFITS VERIFIED

PA STATUS / THROUGH

SCHEDULED



GREATER BOSTON INFUSION CENTER

17/19 Kendrick St, Wrentham, MA 02093
P 774-581-0032 | F 774-581-0033

PAGE 2

Prescribing Order - Benlysta (belimumab)

☐ New Start

☐ Maintenance

DATE OF LAST DOSE / INFUSION

ORDER DETAILS

DATE OF ORDER

PATIENT NAME

DOB

M/F

DIAGNOSIS / ICD-10

☐ NKDA

PATIENT WEIGHT

ALLERGIES

PREMEDICATIONS

☐ Acetaminophen 1000mg PO

☐ Diphenhydramine 25mg IV

☐ Loratadine 10mg or Cetirizine 10mg PO

☐ Solu-medrol 125mg IV

☐ Diphenhydramine 25mg PO

☐ Other

LAB ORDERS

☐



GREATER BOSTON INFUSION CENTER

17/19 Kendrick St, Wrentham, MA 02093
P 774-581-0032 | F 774-581-0033

PAGE 3

Prescribing Order - Benlysta (belimumab) (continued)

BENLYSTA MEDICATION ORDER

- ☐ Loading: 10mg/kg IV at week 0, 2, 4
- ☐ Maintenance: 10mg/kg IV every 4 weeks
- ☐ Other

ADMINISTRATION & MONITORING

- ✓ Do not administer if patient has active signs or symptoms of infection.
- ✓ Do not administer if patient is suspected to have or experiencing symptoms of PML
- ✓ Reconstitute Benlysta with sterile water per manufacturer instructions and dilute in 250ml Normal Saline.
- ✓ Administer over 1 hour.
- ✓ In case of infusion reaction, STOP infusion and follow GBIC infusion reaction protocol. Notify physician.
- ✓ Signature Date

PROVIDER AUTHORIZATION

ORDERING PROVIDER NAME

NPI

DATE

SIGNATURE

OFFICE PHONE / EXT.

Handwritten or compliant electronic signature