



GREATER BOSTON INFUSION CENTER

17/19 Kendrick St, Wrentham, MA 02093
P 774-581-0032 | F 774-581-0033

PAGE 1

Simponi Aria - Referral Checklist

Referral packet checklist

Complete this page and attach the supporting items below.

PATIENT & REFERRING PROVIDER

PATIENT NAME

DOB

DATE

REFERRING PROVIDER

NPI

DIRECT CONTACT / EXT.

OFFICE / PRACTICE

PHONE

FAX

EMAIL

URGENT CLINICAL CONTACT

REQUIRED REFERRAL PACKET

- ☐ Patient demographics and contact information
- ☐ Primary and secondary insurance information; front/back card copies
- ☐ Supporting clinical notes / office visits documenting Simponi Aria treatment plan
- ☐ Current medication list and allergies
- ☐ Relevant laboratory results / imaging and baseline safety information
- ☐ Completed Simponi Aria prescribing order (following page)

MEDICATION-SPECIFIC CLINICAL DOCUMENTATION

- Note should include any therapies tried/failed, and must include discussion about Simponi Aria
- Medication list and allergies
- Required Baseline: TB screening, HBV screening, CBC w diff
- Recommended: periodic CBC w diff

GBIC INTERNAL USE

REFERRAL RECEIVED

BENEFITS VERIFIED

PA STATUS / THROUGH

SCHEDULED



GREATER BOSTON INFUSION CENTER

17/19 Kendrick St, Wrentham, MA 02093

P 774-581-0032 | F 774-581-0033

PAGE 2

Prescribing Order - Simponi Aria (golimumab)

☐ New Start

☐ Maintenance

DATE OF LAST DOSE / INFUSION

ORDER DETAILS

DATE OF ORDER

PATIENT NAME

DOB

M/F

DIAGNOSIS / ICD-10

☐ NKDA

PATIENT WEIGHT

ALLERGIES

PREMEDICATIONS

☐ Acetaminophen 1000mg PO

☐ Diphenhydramine 25mg IV

☐ Loratadine 10mg or Cetirizine 10mg PO

☐ Solu-medrol 125mg IV

☐ Diphenhydramine 25mg PO

☐ Other

LAB ORDERS

☐ CBC w/diff every

☐ CMP every

☐ Other



GREATER BOSTON INFUSION CENTER

17/19 Kendrick St, Wrentham, MA 02093
P 774-581-0032 | F 774-581-0033

Prescribing Order - Simponi Aria (golimumab) (continued)

ENTYVIO MEDICATION ORDER

Dosing: ☐ Simponi Aria 2mg/kg IV

Frequency: ☐ Dose at week 0, 4 and 8 ☐ Maintenance dose every 8 weeks thereafter

ADMINISTRATION & MONITORING

- ✓ Dilute Simponi Aria in 100ml 0.9% Sodium Chloride. Administer intravenously over 30 min
- ✓ Do not administer if patient has active signs or symptoms of infection.
- ✓ Do not administer in patient with signs/symptoms of CHF, TB, or hepatitis
- ✓ In case of infusion reaction, STOP infusion and follow GBIC infusion reaction protocol. Notify physician.
- ✓ Signature Date

PROVIDER AUTHORIZATION

ORDERING PROVIDER NAME	NPI	DATE
<input type="text"/>	<input type="text"/>	<input type="text"/>

SIGNATURE

OFFICE PHONE / EXT.

Handwritten or compliant electronic signature