



GREATER BOSTON INFUSION CENTER

17/19 Kendrick St, Wrentham, MA 02093

P 774-581-0032 | F 774-581-0033

PAGE 1

Tysabri - Referral Checklist

Referral packet checklist

Complete this page and attach the supporting items below.

PATIENT & REFERRING PROVIDER

PATIENT NAME

DOB

DATE

REFERRING PROVIDER

NPI

DIRECT CONTACT / EXT.

OFFICE / PRACTICE

PHONE

FAX

EMAIL

URGENT CLINICAL CONTACT

REQUIRED REFERRAL PACKET

- ☐ Patient demographics and contact information
- ☐ Primary and secondary insurance information; front/back card copies
- ☐ Supporting clinical notes / office visits documenting Tysabri treatment plan
- ☐ Current medication list and allergies
- ☐ Relevant laboratory results / imaging and baseline safety information
- ☐ Completed Tysabri prescribing order (following page)

MEDICATION-SPECIFIC CLINICAL DOCUMENTATION

- Note should include any therapies tried/failed, and must include discussion about Tysabri
- Medication list and allergies
- Baseline: MRI Brain, JCV antibody, CBC w diff, CMP, Varicella antibody
- Recommended for continued therapy: frequent MRI Brain, JCV, CBC w diff, and CMP monitoring

GBIC INTERNAL USE

REFERRAL RECEIVED

BENEFITS VERIFIED

PA STATUS / THROUGH

SCHEDULED



GREATER BOSTON INFUSION CENTER

17/19 Kendrick St, Wrentham, MA 02093
P 774-581-0032 | F 774-581-0033

PAGE 2

Prescribing Order - Tysabri (natalizumab)

☐ New Start

☐ Maintenance

DATE OF LAST DOSE / INFUSION

ORDER DETAILS

DATE OF ORDER

PATIENT NAME

DOB

M/F

DIAGNOSIS / ICD-10

☐ NKDA

PATIENT WEIGHT

ALLERGIES

PREMEDICATIONS

☐ Acetaminophen 1000mg PO

☐ Diphenhydramine 25mg IV

☐ Loratadine 10mg or Cetirizine 10mg PO

☐ Solu-medrol 125mg IV

☐ Diphenhydramine 25mg PO

☐ Other

LAB ORDERS

☐ JCV antibody every

☐ CBC w diff, CMP every



GREATER BOSTON INFUSION CENTER

17/19 Kendrick St, Wrentham, MA 02093
P 774-581-0032 | F 774-581-0033

PAGE 3

Prescribing Order - Tysabri (natalizumab) (continued)

TYSABRI MEDICATION ORDER

- ☐ Tysabri 300mg in 100ml NS IV over 1 hour every 4 weeks x 12 months
- ☐ Tysabri 300mg in 100ml NS IV over 1 hour every weeks x 12 months
- ✓ Monitor patient for 1 hour after Tysabri infusion for first 12 treatments, then monitor per GBIC protocol. Infuse

ADMINISTRATION & MONITORING

- ✓ Do not administer if patient has active signs or symptoms of infection.
- ✓ Do not administer if any suspected signs/symptoms of PML.
- ✓ Complete required TOUCH screening and checklist prior to each infusion.
- ✓ In case of infusion reaction, STOP infusion and follow GBIC infusion reaction protocol. Notify physician.
- ✓ Signature Date

PROVIDER AUTHORIZATION

ORDERING PROVIDER NAME

NPI

DATE

SIGNATURE

OFFICE PHONE / EXT.

Handwritten or compliant electronic signature